

Blue Ribbon Bene-lyte

★★★★★★ For sick/stressed or transported calves.

USAGE INSTRUCTIONS

Follow mixing and feeding directions carefully. Consult your veterinarian for advice and treatment of disease.

INGREDIENTS

Lactose, Dextrose, Sodium Bicarbonate, Citric Acid, Sodium Chloride, Tricalcium Phosphate, Corn Starch, Artificial Flavors, Glycine, Potassium Chloride, Guar Gum, Sodium Aluminosilicate, Blue No. 1.

FEATURES

- ★ **Effervescent mixing**
Due to advanced effervescent technology, Bene-lyte mixes easily with water and does not settle out of solution.
- ★ **Blue coloring**
The addition of blue coloring easily indicates to producers which calves have already consumed Bene-lyte.
- ★ **Convenient packaging**
Bene-lyte is packaged in bulk 10 kg (22 lb) pails for easy handling in larger operations and providing per dose savings.

CALF FEEDING SCHEDULE

	For transported calves	For sick/stressed calves	For weaned calves
Day 1	2 qts 1-2 times daily + 2 qts Milk Replacer 2 times daily	2 qts 2 times daily + 2 qts Milk Replacer 2 times daily	2 qts 3-4 times daily
Day 2	2 qts 1-2 times daily + 2 qts Milk Replacer 2 times daily	2 qts 1-2 times daily + 2 qts Milk Replacer 2 times daily	2 qts 2-3 times daily
Day 3	Milk Replacer	2 qts 1-2 times daily + 2 qts Milk Replacer 2 times daily	2 qts 1-2 times daily
Day 4	Milk Replacer	Milk Replacer	2 qts 1 time daily

MIX & FEED INSTRUCTIONS

Mix one scoop [4 oz (113.5 g)] of Bene-lyte to 2 qts (2 L) of 90°-110° F water in a feeding bucket. Allow the mixing action to subside before feeding. To minimize overflowing when bottle feeding, fill bottles 1/2 full before adding Bene-lyte and add remaining water after mixing action subsides. For best results feed 2-4 hours after feeding milk or milk replacer. If you have any questions, consult Merrick's Customer Service at 1-800-MER-RICK (637-7425).

PRODUCT NO. & PACKAGING

35910123.....10 kg Pail